

Feline Rhinotracheitis-Calici-**Panleukopenia Vaccine**

Modified Live Virus

Focus Cat Vax B

Package contains a one dose vial of dry vaccine, a 1 mL vial of liquid diluent, and one disposable syringe

FOR CATS ONLY

PERISHABLE PRODUCT . KEEP REFRIGERATED

Indications: For the vaccination of healthy cats, 8 weeks of age or older, as an aid in the prevention of disease caused by feline rhinotracheitis, calici and panleukopenia viruses.

Directions and Dosage: The dosage is 1 mL injected subcutaneously using aseptic technique. Open syringe by twisting or tapping the cap against a hard surface to break the heat weld. Prepare the vaccine by injecting the diluent into the vial containing the desiccated vaccine cake. Shake well. Withdraw entire contents into the syringe. Subcutaneous administration: Lift the loose skin behind the neck or behind right hind leg (see illustration) and insert needle. Inject entire contents of syringe. Do not inject directly into blood vessel.

Important note: Before injecting vaccine pull back slightly on syringe plunger. If blood enters the syringe freely, choose another injection site. Vaccinate kittens at 8 weeks of age or older. Administer a second 1 mL dose 3-4 weeks later. Annual revaccination is recommended.

Precautions: Store out of direct sunlight at 35°-45°F (2°-7°C). Do not freeze. Do not vaccinate pregnant animals. Only vaccinate healthy animals. Burn containers and unused contents. Although rare, severe allergic reactions (anaphylactoid) may occur that require immediate veterinary care. Antidote: Epinephrine.

Preservatives: Neomycin and Polymyxin B.



Produced by: Boehringer İngelheim Vetmedica, Inc. St. Joseph, MO 64506 U.S.A. U.S. Vet. License No. 124

Distributed by: Peak Marketing P.O. Box 279 Blue Springs, MO 64013 U.S.A.

Focus® Cat Vax 3 is a trademark of Peak Marketing LLC ISS12XB10

Step 1 Transfer liquid contents of

diluent into vial of dried vaccine

Step 2 Shake well



Step 3 Withdraw mixture into

Step 4 To administer subcutaneously, gently grasp and elevate a fold of loose skin

Step 5 Insert the needle into the loose tissue under

the skin

